

Nucleation of H₂SO₄ and oxidized organics in CLOUD experiment

Mikko Sipilä, Nina Sarnela, Tuija Jokinen, Tuomo Nieminen, Joao Almeida, Jasmin Tröstl, Katrianne Lehtipalo, Jonathan Duplissy, Heikki Junninen, and CLOUD Collaboration

Citation: [AIP Conference Proceedings](#) **1527**, 318 (2013); doi: 10.1063/1.4803267

View online: <http://dx.doi.org/10.1063/1.4803267>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1527?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Multi-species nucleation rates in CLOUD](#)

AIP Conf. Proc. **1527**, 326 (2013); 10.1063/1.4803269

[Aerosol nucleation and growth in a mixture of sulfuric acid/alpha-pinene oxidation products at the CERN CLOUD chamber](#)

AIP Conf. Proc. **1527**, 322 (2013); 10.1063/1.4803268

[Molecular steps of neutral sulfuric acid and dimethylamine nucleation in CLOUD](#)

AIP Conf. Proc. **1527**, 302 (2013); 10.1063/1.4803263

[Measuring composition and growth of ion clusters of sulfuric acid, ammonia, amines and oxidized organics as first steps of nucleation in the CLOUD experiment](#)

AIP Conf. Proc. **1527**, 298 (2013); 10.1063/1.4803262

[Cirrus cloud mimics in the laboratory: An infrared spectroscopy study of thin films of mixed ice of water with organic acids and ammonia](#)

J. Chem. Phys. **126**, 084702 (2007); 10.1063/1.2464082

Nucleation of H₂SO₄ and Oxidized Organics in CLOUD experiment

Mikko Sipilä^a, Nina Sarnela^a, Tuija Jokinen^a, Tuomo Nieminen^a, Joao Almeida^b, Jasmin Tröstl^c, Katrianne Lehtipalo^a, Jonathan Duplissy^a, Heikki Junninen^a, and the CLOUD collaboration

^a University of Helsinki, Department of Physics, P.O. Box 64, 00014 University of Helsinki, Finland

^b Institute for Atmospheric and Environmental Sciences, Goethe-University of Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

^c Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, OFLA/110, 5232 Villigen PSI, Switzerland

Abstract. The research of atmospheric new particle formation has proceeded lately as the role of sulphuric acid has been established. Still, the roles of other atmospheric compounds in nucleation remain largely unclear. To clarify the first steps of atmospheric new particle formation extensive nucleation experiments were performed in CLOUD chamber in 2012. Especially the role of oxidations products of α -pinene was studied in detail. The experiments provided new information about the part of oxidized organics in nucleation.

Keywords: nucleation, chamber measurements, atmospheric aerosols

PACS: 82.60.Nh, 82.33.Tb, 92.60.Mt

INTRODUCTION

Detailed steps of atmospheric new particle nucleation have been under debate and subject to numerous experimental and theoretical studies for decades. Despite the serious efforts, the exact molecular steps of nucleation remain unknown, though recent advances in mass-spectrometry^{1,2,3} and particle counting technologies^{4,5} are finally pointing toward a breakthrough. Sulphuric acid, H₂SO₄, has an established role in atmospheric nucleation^{5,6,7,8} but sulphuric acid alone, or with water, cannot account for the high nucleation rates observed in the boundary layer⁹. Therefore, it has been suggested that additional vapours including bases – such as ammonia⁹ or amines^{10,11,12} – or oxidized organics^{13,14} nucleate together with sulphuric acid and help the formation of new particles. Besides ternary vapours, it has been demonstrated that ions, produced in the atmosphere by galactic cosmic radiation and radon decay, can enhance the nucleation rate of new particles at least in sulphuric acid - water - ammonia system⁹ potentially connecting the fluctuations of cosmic radiation to cloudiness and climate variations¹⁵. Some atmospheric observations, however, suggest that such a connection is weak or non-existent¹⁶. Here we report on our findings on the role of oxidized organics in neutral and ion-induced sulphuric acid nucleation in absence and in presence of ammonia and dimethyl amine.

METHODS

Experiments were performed at CLOUD facility⁹ in CERN during the CLOUD 7 run in autumn 2012. In the experiments α -pinene – a monoterpene, representative especially for boreal forests – was oxidized either with OH only or ozone only or with concurrent presence of both oxidants. Experiments were performed either in presence or in absence of ions in the chamber. Ions were produced in the CLOUD chamber by natural galactic cosmic radiation or, at enhanced concentrations by using CERN proton synchrotron pion beam. In the neutral experiments the complete absence of ions was achieved by sweeping the ions out from the chamber by high electric field. Some of the experiments were done in presence of varying concentrations of dimethyl amine or ammonia.

Experiments utilized a suite of recently developed instruments tailored for the purposes of the nucleation studies. Instruments comprise especially a Chemical – Ionization – Atmospheric Pressure Interface Time of Flight mass spectrometer (CI-API-TOF) developed for detection of gas phase sulphuric acid and nucleating sulphuric acid containing clusters, as well as oxidized organic reaction products³, a Particle Size Magnifier⁴ and other instrumentation (e.g. NAIS, DEG-CPC etc.) for detecting the particles immediately after their nucleation. Nucleation studies of sulphuric acid and amines in absence of α -pinene and its products are described elsewhere¹⁷.

The CI-API-TOF^[3] relays on chemical ionization of sample molecules and clusters by nitrate ions. In the new version of the CI-API-TOF, developed for the CLOUD 7 campaign, the primary ions are generated by irradiating nitric acid containing sheath flow by photons from a soft x-ray source. Main advantages of the x-ray source in comparison to Am-241 source used in original instrument³ are that it is non-radioactive and can be switched on and off. Primary ions are then mixed with the sample flow by means of electric fields. CI-API-TOF can be used for detection of strong acids and acidic clusters that can donate a proton to nitrate ion, as well as certain compound groups, including highly oxidized organics, which can form stable ionic clusters with nitrate ions.

RESULTS

The nucleation rates of experiments were calculated and the values were compared with the prevailing conditions. Especially the correlation with sulphuric acid and oxidized organics was studied in detail. The concentration of sulphuric acid and oxidized organics were calculated from the CI-API-TOF data. The monomers of oxidized organics were considered to be organic compounds between mass charge values from 300 to 450. Figure 1 shows an example of particle formation in an experiment with α -pinene, SO₂ and dimethyl amine. The effects of different concentrations of oxidized organics and sulphuric acid on nucleation rates in presence and in absence of dimethyl amine or ammonia, were studied and the results will be discussed.




Figure 1. An example of the behaviour of particles measured by NAIS and sulphuric acid and oxidized organics monomers measured by CI-API-TOF during an experiment where α -pinene and dimethyl amine was added into the chamber.

ACKNOWLEDGMENTS

We would like to thank CERN for supporting CLOUD with important technical and financial resources, and for providing a particle beam from the CERN Proton Synchrotron. This research has received funding from the EC Seventh Framework Programme (Marie Curie Initial Training Network "CLOUD-ITN" no. 215072, MC-ITN "CLOUD-TRAIN" no. 316662, and ERC-Advanced "ATMNUCLE" grant no.

227463), the German Federal Ministry of Education and Research (project nos. 01LK0902A and 01LK1222A), the Swiss National Science Foundation (project nos. 200020_135307 and 206620_130527), the Academy of Finland (Center of Excellence project no. 1118615), the Academy of Finland (135054, 133872, 251427, 139656, 139995, 137749, 141217, 141451), the Finnish Funding Agency for Technology and Innovation, the Nessling Foundation, the Austrian Science Fund (FWF; project no. P19546 and L593), the Portuguese Foundation for Science and Technology (project no. CERN/FP/116387/2010), the Swedish Research Council, Vetenskapsrådet (grant 2011-5120), the Presidium of the Russian Academy of Sciences and Russian Foundation for Basic Research (grants 08-02-91006-CERN and 12-02-91522-CERN), and the U.S. National Science Foundation (grants AGS1136479 and CHE1012293).

REFERENCES

1. M. Ehn, E. Kleist, H. Junninen, T. Petäjä, G. Lönn, S. Schobesberger, M. Dal Maso, A. Trimborn, M. Kulmala, D. R. Worsnop, A. Wahner, J. Wildt, and Th. F. Mentel, *Atmos. Chem. Phys.*, **12**, 5113-5127 (2012).
2. J. Jiang, J. Zhao, M. Chen, F. L. Eisele, J. Scheckman, B. J. Williams, C. Kuang and P. H. McMurry, *Aerosol Sci. Technol.*, **45**, ii — v (2011)
3. T. Jokinen, M. Sipilä, H. Junninen, M. Ehn, G. Lönn, J. Hakala, T. Petäjä, R. L. Mauldin III, M. Kulmala and D. R. Worsnop, *Atmos. Chem. Phys.* **12**, 4117-4125 (2012)
4. J. Vanhanen, J. Mikkilä, K. Lehtipalo, M. Sipilä, H. E. Manninen, E. Siivola, T. Petäjä and M. Kulmala, *Aerosol Sci. Technol.* **45**, 533-542 (2011)
5. C. Kuang, M. Chen, J. Zhao, J. Smith, P. H. McMurry, and J. Wang, *Atmos. Chem. Phys.*, **12**, 3573–3589 (2012)
6. R. J. Weber, J. J. Marti, P. H. McMurry, F. L. Eisele, D. J. Tanner and A. Jefferson, *Chem. Eng. Comm.* **151**, 53-64 (1996)
7. R. Zhang, *Science*, **11**, 1366-1367 (2010)
8. M. Sipilä, T. Berndt, T. Petäjä, D. Brus, J. Vanhanen, F. Stratmann, J. Patokoski, R. L. Mauldin III, A.-P. Hyvärinen, H. Lihavainen and M. Kulmala, *Science*, **327**, 1243-1246 (2010)
9. J. Kirkby et al, *Nature*, **476**, 429-433 (2011)
10. T. Berndt, T. F. Stratmann, M. Sipilä, J. Vanhanen, T. Petäjä, J. Mikkilä, A. Grüner, G. Spindler, R. L. Mauldin III, J. Curtius, M. Kulmala, and J. Heintzenberg, *Atmos. Chem. Phys.* **10**, 7101-7116 (2010).
11. J. Zhao, J. N. Smith, F. L. Eisele, M. Chen, C. Kuang, and P. H. McMurry, *Atmos. Chem. Phys.*, **11**, 10823-10836 (2011)
12. M. Chen, M. Titcombe, J. Jiang, C. Jen, C. Kuang, M. L. Fischer, F. L. Eisele, J. I. Siepmann, D. R. Hanson, J. Zhao and P. H. McMurry, *Proc. Nat. Acad. Sci.*, **109**, 18713–18718 (2012)
13. A. Metzger, B. Verheggen, J. Dommen, J. Duplissy, A. S. Prevot, E. Weingartner, I. Riipinen, M. Kulmala, D. V. Spracklen, K. S. Carslaw and U. Baltensperger, *Proc. Nat. Acad. Sci.*, **107**, 6646-6651 (2010)
14. F. Riccobono, L. Rondo, M. Sipilä, P. Barmet, J. Curtius, J. Dommen, M. Ehn, S. Ehrhart, M. Kulmala, A. Kürten, J. Mikkilä, P. Paasonen, T. Petäjä, E. Weingartner, and U. Baltensperger, *Atmos. Chem. Phys.*, **12**, 9427-9439 (2012).
15. H. Svensmark and E. Friis-Christensen, *J. Atmos. Terr. Phys.*, **59**, 1224–1232 (1997)
16. M. Kulmala, I. Riipinen, T. Nieminen, M. Hultkonen, L. Sogacheva, H. E. Manninen, S. Paasonen, P., Petäjä, T., Dal Maso, M., Aalto, P. P., Viljanen, A., Usoskin, I., Vainio, R., Mirme, A. Mirme, A. Minikin, A. Petzold, U. Hörrak, C. Plaß-Dülmer, W. Birmili and V.-M. Kerminen, *Atmos. Chem. Phys.*, **10**, 1885-1898 (2010)
17. Jokinen et al., 2013, these proceedings